AD		

Award Number: DAMD17-99-1-9267

TITLE: Long Interspersed Nuclear Element (LINE) Mediated Genomic Instability

PRINCIPAL INVESTIGATOR: Lalitha Nagarajan, Ph.D.

CONTRACTING ORGANIZATION: The University of Texas
MD Anderson Cancer Center
Houston, Texas 77030

REPORT DATE: July 2000

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

### REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

3. REPORT TYPE AND DATES COVERED 1. AGENCY USE ONLY (Leave 2. REPORT DATE blank) July 2000 Annual (1 Jul 99 - 30 Jun 00) 4. TITLE AND SUBTITLE 5. FUNDING NUMBERS DAMD17-99-1-9267 Long Interspersed Nuclear Element (LINE) Mediated Genomic Instability 6. AUTHOR(S) Lalitha Nagarajan, Ph.D. 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER The University of Texas MD Anderson Cancer Center Houston, Texas 77030 E-MAIL: lnagarajan@notes.mdacc.tmc.edu 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING / MONITORING AGENCY REPORT NUMBER U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 11. SUPPLEMENTARY NOTES 12a, DISTRIBUTION / AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE Approved for public release; distribution unlimited

### 13. ABSTRACT (Maximum 200 Words)

We have isolated and characterized a novel human homeobox gene designated Mix.1 homeobox (Xenopus laevis)-like gene 2 (MIXL2), as a gene that contains a repetitive element in its 3'UTR (MER22). The MIXL2 gene encodes a predicted 232 amino acid open reading frame, which contains a highly conserved mix.1-like homeobox and an acidic motif. Radiation hybrid mapping and Fluorescence in situ hybridization localize the MIXL2 gene to human chromosome 1q32-41, a region of gain in breast tumors. Consistent with these cytogenetic findings we detected allelic imbalance for loci flanking MIXL2 in 14/25 primary breast tumors. Additionally, a search of the human dbEST database suggested a highly restricted expression as we detected only three EST clones (one each from normal germinal center B cell, undifferentiated germ cell tumor, and infiltrating ductal carcinoma of the breast). These findings suggest MIXL2 may play a role in development and mammary carcinogenesis.

14. SUBJECT TERMS Breast Cancer, geneomi homeobox gene	c insatbility, chromoso	omal gain, 1q41 and	15. NUMBER OF PAGES 16 16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Inclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	Unlimited

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. Z39-18 298-102

### FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

N/A Where copyrighted material is quoted, permission has been obtained to use such material.

N/A Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

N/A Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

 $\frac{N/A}{L}$  In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

 $\underline{N/A}$  In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

 ${\rm N/A}$  In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

La litha not nage 7/19/00 PI - Signature Date

### **Table of Contents**

Cover1	
SF 2982	2
Foreword	3
Table of Contents4	ļ
Introduction	.5
Body	6-7
Key Research Accomplishments8	
Reportable Outcomes9	ı
Conclusions10	0
References11	I
Appendices12	2-16

### Introduction:

The overall goals of this proposal are to develop a rapid PCR assay to detect the genome wide Long Interspersed Sequence (L1Hs) mediated instability and to isolate target oncogenes, tumor suppressor genes and regions of chromosomal instability by the use of L1Hs specific PCR. In order to accomplish these goals we modified our approach and took advantage of the recent progress in the human genome intiative and cancer genome anatomy projects (CGAP).

DNA sequence analysis of a novel L1Hs element that we characterized in the laboratory revealed a segment of homology to MER22 repetitive elements. These findings coupled with the rapid progress in the human genome initiative led us to redesign our approach to detect target genes. Thus we screened the cDNA database dBEST for novel genes which contain repetitive elements in them. About the same time, other studies in our laboratory identified a partial cDNA clone for a key regulatory homeobox MIX like gene in the dBEST database. Interestingly, the 3' UTR of this gene contains MER repetitive elements. Furthermore the expression of this novel gene appeared to be highly restricted with the identification only 3 cDNA clones in the entire dBEST database with more than 3 million entries. Our interest in this regulatory gene was further heightened as one of the 3 cDNA clones was from a laser captured tumor tissue from infiltrating mammary adenocarcinoma *in situ*. All of the subsequent studies detailed in this report stemmed from these key observations made in early stages of the first year of this award.

Year 1 of the proposal was to focus on specific Aim 1. The following report summarizes what we have accomplished towards specific tasks in the approved statement of work.

### Specific Aim I (i-iii)

- (i) Development of primers and conditions for IPCR and L1Hs PCR
- (ii) Analysis of hybridization patterns
- (iii) Subcloning of unique fragments

In accomplishing the tasks i-iii of Specific Aim I, we have taken advantage of the rapid progress made in the human genome sequencing initiative, the cancer genome anatomy project and more importantly characterization of a novel L1Hs element in our laboratory. The novel element designated LREHL showed segments of homology to MER repetitive elements.

We initiated a screen of the expressed sequence tagged database (dBEST) electronically for transcripts with repetitive sequences. About the same time, other investigations in our laboratory were searching for the human homolog of the homeobox gene *Mix.1*, which regulates mesoderm and endoderm differentiation in Xenopus embryos. A single candidate EST clone (AA84781) with a putative *Mix* like homeodomain was identified in a library from germinal center B lymphoid cells. When the EST clone AA84781 was sequenced, we identified a partial open reading frame. A recursive search of the dBEST database identified only two other cDNA clones for this novel gene, suggesting an extremely restricted pattern of expression. Interestingly, one of these ESTs (AA911377) was from a cDNA library made from laser captured infiltrating ductal carcinoma of the breast (Fig.1). The significance of this finding was further enhanced when we detected alu and MER like repetitive elements in the 3' (> 1.0kbp) UTR of this gene.

### (iv) Sequence analysis and development of STSs

We developed unique STSs from the 3'untranslated region of the MIX like homeodomain gene. Using these primers we screened a human genomic library in Bacterial Artificial Chromosome vector to isolate a large segment of the genome containing the gene. The BAC was used to characterize the full length open reading frame of this novel homeogene that we designated *MIXL2*. The gene encodes an open reading frame of 232amino acids. Similar genes whose expression is restricted to the primitive streak have been identified in mice and chicken (1-3). MIXL2 shows 94% identity in mix-like domain with an overall identity of 69% to the murine gene MML (Fig.2). With the chicken gene CMIX, we find a 79% identity in the homeodomain and an overall identity of 41%. Identification of the three distinct and yet highly related genes suggests the existence of a novel mix-like homeobox gene family.

Additionally, we characterized the genomic organization and expression pattern of MIXL2. The gene consists of two exons (Fig. 3) and shows a highly restricted expression pattern. None of the breast cancer cell lines tested showed abundant transcripts. However, reverse transcription coupled PCR detected *MIXL2* transcripts in the cell line MDA-MB 453 and not in the immortalized breast cell line MCF10A.

Simultaneously, we fine mapped the novel gene to human genome using DNA from a panel of well-characterized radiation hybrids. Interestingly, the MIX like gene localized to human chromosome 1q32-41 locus, between the microsatellite loci *D1S479* and, a region of gain in breast cancer (Fig.4).

### (v) Development of the PCR for minimal material

In collaboration with Dr. Nour Sneige in the Dept. of Pathology, we dissected 25 breast tumors and isolated normal and tumor DNAs. The status of D1S479 was examined in these paired samples by polymerase chain reaction (PCR) with end labeled primers. Interestingly, 12 out of 25 tumors showed allelic imbalance for *D1S479* suggesting that the MIXL gene was a candidate target gene of chromosomal gain in 1q (Fig. 5).

### (vi) Correlation of data from tasks (i-v) with stage and grade

All of the 25 breast tumors used to screen for allelic imbalance of *D1S479* showed lymph node involvement. Studies are underway to determine whether there is intra-tumoral heterogeneity within for gain of 1q32-41 loci. We will utilize both FISH and MIXL specific antibody based immunofluorescence staining approaches to examine such a possibility.

### **Key Research Accomplishments:**

- Identification of a novel regulatory homeobox gene with repetitive elements in the 3'UTR
- Isolation and characterization of the full length open reading frame for the novel gene designated *MIXL2*
- Localization of MIXL2 to 1q41, a region of gain in breast cancer
- Detection of allelic imbalance for *D1S479*, a marker physically linked to *MIXL2*, in 12/25 breast tumors with lymph node involvement

### Reportable outcomes:

Manuscripts in preparation:

- 1. Hejlik DP and Nagarajan L. Retrotransposition and recombination of LINE-1 Elements as a mechanism of chromosomal deletion and genomic instability
- 2. Guo W., Ma, J., Liang H., and Nagarajan L. A novel MIX like paired type homeobox gene localizes to a region of in breast cancer
- 3. Guo W., Chan A., Ma J., Liang H., Etkin L and Nagarajan L. MIXL2 shows evolutionary and functional conservation: a role in hematopoiesis

### Development of FISH and immunohistochemical staining probes:

We have characterized a BAC probe that can be used in Fluorescence in situ hybridization (FISH).

We have developed antibodies to MIXL2, which can be used in immunohistochemical staining.

### Informatics:

The cDNA sequence for MIXL2 has been submitted to human genome organization. The full length sequence will be submitted to Genbank.

### Funding applied for based on work supported by this award:

Application by Wei Guo entitled "Functional characterization of a novel embryonic homeobox gene MIXL2 in breast cancer" for Graduate training fellowship, from the Breast Cancer Research Program, Department of Defense Submitted on June 7, 2000.

### **Conclusions:**

A genome wide-electronic screening approach yielded us clues on an important regulatory homeobox gene with repetitive elements in the 3' UTR.

The MIXL2 gene is highly restricted in its expression. There are only 3 cDNA clones for this gene in the entire human cDNA expressed sequence tag (EST) database that contains more than 3 million entries. Interestingly, one of the three ESTs for MIXL2 is from a laser captured infiltrating ductal carcinoma in situ of the breast.

The *MIXL2* gene consists of two exons and encodes an open reading frame of 232 amino acids.

We localized it to a region of chromosomal gain (1q41) in breast cancer.

Interestingly *D1S479*, a marker that is physically linked to *MIXL2* is gained in 12 out of 25 breast tumors with lymph node involvement.


Homeobox genes encode a family of regulatory transcription factors that primarily play a crucial role in development. Other evidences suggest that certain homeobox genes provide an important link between the processes of cell cycle control, development and cancer.

Future experiments will identify the functional consequence of gain of *MIXL2* gene in breast cancer as well as characterization of other novel regulatory genes with L1 repetitive elements.

### References:

- 1. Pearce JJ, Evans MJ. Mml, a mouse Mix-like gene expressed in the primitive streak. *Mech. Dev.* **87**:189-192,1999.
- 2. Stein S, Roeser T, Kessel M. CMIX, a paired-type homeobox gene expressed before and during formation of the avian primitive streak. *Mech. Dev.* **75**:163-165,1998.
- 3. Peale FV Jr, Sugden L, Bothwell M. Characterization of CMIX, a chicken homeobox gene related to the Xenopus gene mix.1. *Mech. Dev.* **75**: 167-170,1998.

### dbEST Database Search



EST	Library	Source
\A847809	NCI_CGAP_GCB1	germinal center B cell
A1654861	NCI_CGAP_GC6	pooled germ cell tumors
AA911377	NCI CGAP Br5	infiltrating ductal carcinoma of the breast


EST clone shows a mix-like homeodomain. The complete sequencing of the EST clone allows us to retrieve the other two EST clones from different cDNA libraries. The detection of only three EST clones suggests a highly restricted expression Figure 1. A search of dbEST database. AA847809 was found in the dbEST database because the partial sequence of the of MIXL2.

# Homology between human MIXL2 and mouse MML

MIXL2	MATAESRALQFAEGAAFPAYRAPHAGGALLPPPSPAAALLPAPPAGPGPATFAGFLGRDP 60
MML	MAAAGSQQLQFAEGAAFPIFPAAHPGGQLLPAMRPASGLPAAPHDSRAPAATQCFPNRDS 60
	** * ** * ** * ** * * * * * * * * * * *
MIXL2	GPAPPPPASLGSPAPPKGAAAPSASQRRKRTSFSAEQLQLLELVFRRTRYPDIHLRERLA 120
MML	SPTAQTPAGLDPPGPSKGSAAPSAPQRRKRTSFSSEQLQLLELVFRQTMYPDIHLRERLA 120
	******** * * * * * * * * * * * * * * * *
MIXL2	ALTLLPESRIQVWFQNRRAKSRRQSGKSFQPLARPEIILNHC-APGTETKCLKPQLPLEV 179
MML	ALTLLPESRIQVWFQNRRAKSRRQSGKSFQPLSSRRGVFLHCPAPGTEARCLKPQLPLEA 180
	**********
MIXL2	DVNCLPEPNGVGGGISDSSSQGQNFETCSPLSEDIGSKLDSWEEHIFSAFGNF 232
MML	DVNHVPDPSMTGGGVCTSGSQSFETYSSLSEDIGSKLDSWEEHIFSALGNF 231
	*** *********** * *** * * * * * * * * *


Figure 2. Homology between human MIXL2 and mouse MML. MML is a mix-like gene in mice. MIXL2 shows 94% identity on the mix-like domain and overall identity of 69% with mouse MML. The carboxy terminal acidic motif is also conserved.

## Genomic Organization of MIXL2



from BAC216I11 sequence. The context around the predicted ATG start codon shows G-3 and G+4 (GGAGCGatgG), which uses a Qt-adapted single strand cDNA pool made from KG1 cells (Methods in Enzym., vol 218, 340-357) to amplify MIXL2 Figure 3. Genomic organization of MIXL2. The predicted MIXL2 consists of at least two exons. The exon1 is predicted to intron junction in the mix-like domain perfectly matches the consensus GT...AG. The part of exon1 sequence is assembled have more than 393bp. 3'-RACE shows that the exon2 has about 1.3kb. The long PCR amplified a 1.4kb intron. The exonmatches the consensus motif of start codon. The EST clone AA847809 covers part of exon1 and exon2. Notes: 3'-RACE exon2 and 3' UTR. The long PCR uses Expand High Fidelity PCR System from Boehringer, and PCR conditions follow

## Genomic Localization of MIXL2



(data not shown). PCR confirms that MIXL2 can hit YAC750G7 in the contig Whitehead Institute Contig WC-516 of Chr1q. The BAC216I11 was retrieved from human BAC library (Research Genetics, Inc) by PCR screening. One end of BAC216I11 also hit Figure 4. Genomic localization of MIXL2.MIXL2 is localized to human chromosome 1q32-41 by radiation mapping and FISH

# Allelic Imbalance of Chromosome 1q32-41in


### Breast Cancer

Allelic imbalance in 12/25 cases

Ġ

Case #	1	7	3	4	\$	9	^	<b>∞</b>	6	10	11	12	13
Accessio	-26SS	-76SS	-26SS	-26SS	-76SS	-76SS	SS97-						
#u	17729	18252	18659	19607	22197	22332	25357	25363	25832	30517	31380	31390	35459
Tumor	+/+	Z	+/+++	Z	N	+/++	IN	Z	Z	++/+	# / +	+/#	##/+
Normal	Z	Z	Z	Z	Z	Z	Z	Z	Z	z	Z	Z	Z

Case #	14	15	16	17	18	19	70	21	22	23	24	25
Accessio	-26SS	-26SS	-26SS	-26SS	-26SS	-26SS	-79SS	-26SS	-26SS	SS97-	-76SS	-79SS
#4	38032	38312	38569	39449	40702	42153	42385	43544	44939	45814	46375	51787
Tumor		+++/+ +/+++	+++/+	##/#	N?	'n.	Z	N?	Z	Z	++/+	+/+
Normal	N	Z	Z	Z	Z	Z	Z	Z	Z	Z	Z	z



œ.